

# 01 Fractions

## CHAPTER LESSON PLANS & OBJECTIVES

### Lesson 1.1: Changing a Fraction

1. Changing an improper fraction to a mixed number
2. Changing a mixed number to an improper fraction
3. Changing a fraction to an equivalent fraction with the lowest common denominator
4. Changing a mixed number to an equivalent fraction with the lowest common denominator

### Lesson 1.2: Adding, Subtracting, Multiplying, Dividing, and Reducing a Fraction

5. Adding fractions having the same denominator, having unlike denominators, or involving whole numbers and unlike denominators
6. Subtracting fractions having the same denominator, having unlike denominators, or involving whole numbers and unlike denominators
7. Multiplying fractions and mixed numbers
8. Dividing fractions and mixed numbers
9. Reducing a complex fraction
10. Reducing a complex fraction involving mixed numbers

# Classroom Preparation

## Lesson 1.1: Changing a Fraction

### INSTRUCTOR PREPARATION

#### Textbook Objectives Covered

1. Changing an improper fraction to a mixed number
2. Changing a mixed number to an improper fraction
3. Changing a fraction to an equivalent fraction with the lowest common denominator
4. Changing a mixed number to an equivalent fraction with the lowest common denominator

# Classroom Preparation

## Lesson 1.1: Changing a Fraction

### STUDENT PREPARATION (3 hrs)

<b>1</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>Changing an Improper Fraction to a Mixed Number</li> </ul>
<b>2</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>Changing a Mixed Number to an Improper Fraction</li> </ul>
<b>3</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>Lowest Common Denominator</li> <li>Changing a Fraction to an Equivalent Fraction with the Lowest Common Denominator</li> </ul>
<b>4</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>Lowest Common Denominator</li> <li>Changing a Mixed Number to an Equivalent Fraction with the Lowest Common Denominator</li> </ul>

# 50-Minute Lesson Plan

## Lesson 1.1: Changing a Fraction

### LECTURE OUTLINE (35 min)

	POWERPOINT SLIDES	CHAPTER HEADING(S)
1	FRACTIONS AND MIXED NUMBERS: SLIDES 3-4	<ul style="list-style-type: none"> <li>Changing an Improper Fraction to a Mixed Number</li> </ul>
2	CHANGING A MIXED NUMBER TO AN IMPROPER FRACTION: SLIDE 5	<ul style="list-style-type: none"> <li>Changing a Mixed Number to an Improper Fraction</li> </ul>
3	CHANGING A FRACTION TO AN EQUIVALENT FRACTION WITH THE LOWEST COMMON DENOMINATOR: SLIDE 6	<ul style="list-style-type: none"> <li>Lowest Common Denominator</li> <li>Changing a Fraction to an Equivalent Fraction with the Lowest Common Denominator</li> </ul>
4	CHANGING A MIXED NUMBER TO AN EQUIVALENT FRACTION WITH THE LOWEST COMMON DENOMINATOR: SLIDE 7	<ul style="list-style-type: none"> <li>Lowest Common Denominator</li> <li>Changing a Mixed Number to an Equivalent Fraction with the Lowest Common Denominator</li> </ul>

### LEARNING ACTIVITIES (choose one or more to equal 15 min)

1	<b>REVIEW (15 min)</b> <ul style="list-style-type: none"> <li>Have the class complete the Pretest. Collect their work and use the results to guide how you teach the class.               <ul style="list-style-type: none"> <li><i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
2	<b>REVIEW (15 min)</b> <ul style="list-style-type: none"> <li>Create a fractions worksheet handout and have students work on it together in groups of three or four. For all Class Activities, periodically rotate through the groups to check on progress and answer questions. Present the answers later to the class as a whole.               <ul style="list-style-type: none"> <li><i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
3	<b>REVIEW (15 min)</b> <ul style="list-style-type: none"> <li>Have two teams of students race at the blackboard to convert fractions to equivalent fractions with the LCD. The team that correctly converts a fraction correctly gets 1 point. The team with the most points wins.               <ul style="list-style-type: none"> <li><i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
4	<b>DISCUSS (15 min)</b> <ul style="list-style-type: none"> <li>Have two teams of students race at the blackboard to convert mixed numbers to equivalent fractions with the LCD. The team that converts a mixed number correctly gets 1 point. The team with the most points wins.               <ul style="list-style-type: none"> <li><i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>

# Classroom Preparation

## Lesson 1.2: Adding, Subtracting, Multiplying, Dividing, and Reducing a Fraction

### INSTRUCTOR PREPARATION

#### Textbook Objectives Covered

##### Lesson 1.2: Adding, Subtracting, Multiplying, Dividing, and Reducing a Fraction

5. Adding fractions having the same denominator, having unlike denominators, or involving whole numbers and unlike denominators
6. Subtracting fractions having the same denominator, having unlike denominators, or involving whole numbers and unlike denominators
7. Multiplying fractions and mixed numbers
8. Dividing fractions and mixed numbers
9. Reducing a complex fraction
10. Reducing a complex fraction involving mixed numbers

# Classroom Preparation

## Lesson 1.2: Adding, Subtracting, Multiplying, Dividing, and Reducing a Fraction

### STUDENT PREPARATION (3 hrs)

<b>5</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Addition of Fractions</li> <li>• Addition of Fractions with the Same Denominator</li> <li>• Addition of Fractions with Unlike Denominators</li> <li>• Addition of Fractions Involving Whole Numbers and Unlike Denominators</li> </ul>
<b>6</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Subtraction of Fractions</li> <li>• Subtraction of Fractions with the Same Denominator</li> <li>• Subtraction of Fractions with Unlike Denominators</li> <li>• Subtraction of Fractions Involving Whole Numbers and Unlike Denominators</li> </ul>
<b>7</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Multiplication of Fractions</li> <li>• Multiplication of Mixed Numbers</li> </ul>
<b>8</b>	<b>READ –Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Division of Fractions</li> <li>• Division of Mixed Numbers</li> </ul>
<b>9</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Reduction of a Complex Fraction</li> </ul>
<b>10</b>	<b>READ – Chapter Heading(s)</b> <ul style="list-style-type: none"> <li>• Reduction of a Complex Fraction</li> <li>• Reduction of a Complex Fraction with Mixed Numbers</li> </ul>

# 50-Minute Lesson Plan

## Lesson 1.2: Adding, Subtracting, Multiplying, Dividing, and Reducing a Fraction

### LECTURE OUTLINE (20 min)

	POWERPOINT SLIDES	CHAPTER HEADING(S)
5	ADDITION OF FRACTIONS: SLIDES 8-11	<ul style="list-style-type: none"> <li>• Addition of Fractions</li> <li>• Addition of Fractions with the Same Denominator</li> <li>• Addition of Fractions with Unlike Denominators</li> <li>• Addition of Fractions Involving Whole Numbers and Unlike Denominators</li> </ul>
6	SUBTRACTION OF FRACTIONS: SLIDES 12-14	<ul style="list-style-type: none"> <li>• Subtraction of Fractions</li> <li>• Subtraction of Fractions with the Same Denominator</li> <li>• Subtraction of Fractions with Unlike Denominators</li> <li>• Subtraction of Fractions Involving Whole Numbers and Unlike Denominators</li> </ul>
7	MULTIPLYING FRACTIONS AND MIXED NUMBERS: SLIDE 15	<ul style="list-style-type: none"> <li>• Multiplication of Fractions</li> <li>• Multiplication of Mixed Numbers</li> </ul>
8	DIVIDING FRACTIONS AND MIXED NUMBERS: SLIDE 16	<ul style="list-style-type: none"> <li>• Division of Fractions</li> <li>• Division of Mixed Numbers</li> </ul>
9	REDUCING A COMPLEX FRACTION: SLIDE 17	<ul style="list-style-type: none"> <li>• Reduction of a Complex Fraction</li> </ul>
10	REDUCING A COMPLEX FRACTION WITH MIXED NUMBERS: SLIDE 18	<ul style="list-style-type: none"> <li>• Reduction of a Complex Fraction</li> <li>• Reduction of a Complex Fraction with Mixed Numbers</li> </ul>

### LEARNING ACTIVITIES (choose one or more to equal 30 min)

5	<b>PRACTICE (15 min)</b> <ul style="list-style-type: none"> <li>• Create a fractions worksheet that requires students to reduce fractions to their lowest terms. Have students work together in groups of three or four to complete it.               <ul style="list-style-type: none"> <li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
6	<b>PRACTICE (20 min)</b> <ul style="list-style-type: none"> <li>• Create a fractions worksheet that requires students to subtract and reduce fractions to the lowest terms. Have students work together in groups of three or four to complete it.               <ul style="list-style-type: none"> <li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
7	<b>PRACTICE (15 min)</b> <ul style="list-style-type: none"> <li>• Create problems involving the multiplication of fractions and mixed numbers, and have two groups of students compete at the board to solve them. The first group to correctly answer the question gets 1 point, and the group with the most points at the end wins.               <ul style="list-style-type: none"> <li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
8	<b>PRACTICE (20 min)</b> <ul style="list-style-type: none"> <li>• Create a fractions and mixed numbers worksheet, and have students work on it together in groups of three or four.               <ul style="list-style-type: none"> <li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li> </ul> </li> </ul>
9	<b>DISCUSS (10 min)</b> <ul style="list-style-type: none"> <li>• Ask students to discuss the relationship among multiplying fractions, dividing fractions, and reducing complex fractions. Then have a volunteer explain how the three processes are</li> </ul>

TEACH Lesson Plan

OGDEN and FLUHARTY: *Calculation of Drug Dosages, 12th Edition*

Copyright © 2022 by Elsevier, Inc. All rights reserved.

	related to one another. Get the class involved and make certain that they understand the relationship before moving on. <ul style="list-style-type: none"><li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li></ul>
<b>10</b>	<b>PRACTICE (15 min)</b> <ul style="list-style-type: none"><li>• Have pairs of students go to the board and compete to reduce complex fractions and to reduce complex fractions containing mixed numbers.<ul style="list-style-type: none"><li>• <i>Appropriate Settings:</i> Traditional classroom, flipped classroom</li></ul></li></ul>



# Assessments

## Chapter 1: Fractions

### ASSESSMENTS BY OBJECTIVE

1	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Fractions Worksheet, Improper Fractions to Mixed Numbers</li> </ul>
2	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Fractions Worksheet, Mixed Numbers to Improper Fractions</li> </ul>
3	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Fractions Worksheet, Add and reduce fractions to lowest terms</li> </ul>
4	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Fractions Worksheet, Mixed Numbers to Improper Fractions</li> </ul>
5	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1: questions 1-7</li> <li>Posttest 2: questions 1-7</li> </ul>
6	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1: questions 8-14</li> <li>Posttest 2: questions 8-14</li> </ul>
7	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1: questions 15-22</li> <li>Posttest 2: questions 15-22</li> </ul>
8	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1, questions 23-30</li> <li>Posttest 2, questions 23-30</li> </ul>
9	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1: questions 1-30</li> <li>Posttest 2: questions 1-30</li> </ul>
10	<b>Study Guide</b> <ul style="list-style-type: none"> <li>Posttest 1: questions 1-30</li> <li>Posttest 2: questions 1-30</li> </ul>

**Objectives:**

- Evaluate orders for minimum and maximum pediatric safe dosage range (SDR) doses
- Performing pediatric dosage calculations
- Converting weight from pounds to kilograms
- Calculate 24-hour drug doses and divided doses for specific weights for oral and parenteral medications

**Case Study:**

M.S. is a 6-month old male that was born at 26 weeks. The infant was discharged 2 weeks ago after a 14 week stay in the neonatal intensive care unit NICU. He currently weighs 6 pounds and is 19 inches long. His current vital signs are HR 145, RR 26, B/P 90/60, T 101.6 F His parents brought the infant to the emergency department because he has not been eating and is continuously fussy. The healthcare provider assessed a bulging left tympanic membrane and diagnosed the infant with an ear infection. Orders include: amoxicillin 113 mg PO q12h. Safe dosing for amoxicillin is 80-90 mg/kg/day divided every 12 hours for 5-7 days.

**Highlight the information that is needed for the nurse to administer the ordered medication safely.**

The nurse is preparing to discharge M.S. **Which nursing action is required to ensure the amoxicillin dose is safe for M.S.? (Select all that apply)**

1. Calculate the infant's weight as 2.7 Kg
2. Calculate the infant's weight as 13.2 Kg
3. Calculate the safe dosage range for amoxicillin as: 216-243 mg/day or 108-121.5 mg/dose every 12 hours
4. Calculate the safe dosage range for amoxicillin as 480-540 mg/day or 240-270 mg/dose every 12 hours

**Choose the *most likely* options for the information missing from the statement below by selecting from the lists of options provided.**

The amoxicillin dose of 113 is \_\_\_\_\_ **1** \_\_\_\_\_ and \_\_\_\_\_ **2** \_\_\_\_\_. The nurse would \_\_\_\_\_ **3** \_\_\_\_\_.

<b>1</b>	<b>2</b>	<b>3</b>
Too high	Safe	Call the physician
Too low	Unsafe	Call the pharmacy
Within range	Lethal	Give the parents the medication

M.S. is brought back to the emergency department by his parents because he appears to have difficulty breathing, 12 hours after being discharged. The parents report this happened after M.S. received his second dose of amoxicillin. The healthcare provider believes M.S. is having an allergic reaction to the

NGN Unfolding case study: Pediatric calculations and safety

amoxicillin. The healthcare provider orders epinephrine 0.01 mg/kg IM now and a normal saline intravenous fluid bolus of 20 mL/kg. The infant weighs 6 pounds.

**Choose the *most likely* option for the missing information from the statements below by selecting from the list of options provided.**

The nurse will administer epinephrine to the infant by \_\_\_\_\_ **1** \_\_\_\_\_ IM into the \_\_\_\_\_ **2** \_\_\_\_\_. The infant will also receive \_\_\_\_\_ **3** \_\_\_\_\_ of normal saline after an intravenous line has been stated.

<b>1</b>	<b>2</b>	<b>3</b>
Using a 1 mL syringe to draw up 0.13 mL of epinephrine 1 mg/mL solution	Vastus Lateralis (Anterolateral thigh)	264 mL
Using a 3 mL syringe to draw up 0.06 mL of epinephrine 1 mg/mL solution	Gluteus maximus	54 mL
Using a 1 mL syringe to draw up 0.03 mL of epinephrine 1 mg/mL solution	Deltoid	20 mL

M.S. was admitted to the hospital for monitoring after receiving the epinephrine and fluid bolus. Current vital signs include: temperature 39 degrees Celsius, HR 150, RR 26, B/P 94/68, sat 99% on room air

The healthcare provider makes morning rounds and orders the following:

Ceftriaxone 50 mg/kg IV daily.

ibuprofen 20 mg PO q8h PRN for temperature greater than 37.5 (include parameter – degrees Celsius)

acetaminophen 64 mg PO q4h PRN for temperature greater than 37.5.

The safe dose range for ibuprofen is 5-10 mg/kg PO q6-8h PRN.

The safe dose range for acetaminophen is 10-15 mg/kg q4-6h PRN.

**Use an “X” (or drag and drop) to indicate which nursing actions listed in the left column would be indicated or contraindicated.**

<b>Actions</b>	<b>Indicated</b>	<b>Contraindicated</b>
Administer Ceftriaxone 300 mg IV		
Administer Ceftriaxone 135 mg IV		
Calculate the safe dosage range for ibuprofen as 13.5-27 mg per dose. The dose of 20 mg PO Q8h PRN is safe and should be administered.		
Calculate the safe dosage range for ibuprofen as 66-132 mg per dose. The dose of 20 mg PO q8H PRN is low and the healthcare provider should be notified.		

NGN Unfolding case study: Pediatric calculations and safety

Calculate the safe dosage range for acetaminophen as 27-40.5 mg per dose. The dosage of 64 mg PO q4h PRN is too high and the healthcare provider should be notified.		
Calculate the safe dosage range for acetaminophen as 60-90 mg per dose. The dosage of 64 mg PO q4h PRN is safe and should be administered.		

Four hours pass after administration of ibuprofen and acetaminophen. The infant currently has a temperature of 38.8 degrees Celsius. Blood pressure one hour ago was 94/62. Current blood pressure is 60/30

M.S. received a fluid bolus of 108 mL of normal saline for a blood pressure of 60/30 and his blood pressure after the fluid bolus is 68/34. The provider suspects early sepsis and orders norepinephrine 0.05 mcg/kg/min and another normal saline fluid bolus of 20 mL/kg.

M.S. weighs 6 pounds. The initial dose of norepinephrine for septic patients is 0.05-0.1 mcg/kg/min. The safe dose for fluid boluses for septic patients is 20-60 mL/kg. The pharmacy provides norepinephrine as 1 mg in 250 mL D5W. Round all values to the nearest tenth. **Use an “X” for the nursing actions listed below that are indicated (necessary), contraindicated (could be harmful), or non-essential (makes no difference or not necessary). Only one selection can be made for each nursing action.**

Nursing Action	Indicated	Contraindicated	Non-Essential
Administer a normal saline fluid bolus of 120 mL			
Administer a normal saline fluid bolus of 54 mL			
Calculate the norepinephrine dose as 1mL/hr			
Calculate the norepinephrine dose as 2.3 mL/hr			
Administer acetaminophen			

After 5 days in the hospital, M.S. has a temperature of 37 degrees Celsius and is off of the norepinephrine drip. The infant’s parents want to take the baby home. The healthcare provider assesses a minor infection remains in the left ear and orders continued antibiotic therapy at home. Orders include

azithromycin 25 mg PO daily for 3 days. M.S. currently weighs 5.5 pounds. The safe dose for azithromycin is 10 mg/kg daily for 3 days. **Highlight the information necessary for the nurse to determine if the azithromycin dose is safe for M.S. before sending him home.**

### Answers and Rationales:

#### Rationale:

##### Phase 1

M.S. is a 6-month old male that was born at 26 weeks. The infant was discharged 2 weeks ago after a 14 week stay in the neonatal intensive care unit NICU. He currently weighs **6 pounds** and is 19 inches long. His current vital signs are HR 145, RR 26, B/P 90/60, T 101.6 F His parents brought the infant to the emergency department because he has not been eating and is continuously fussy. The healthcare provider assessed a bulging left tympanic membrane and diagnosed the infant with an ear infection. Orders include: **amoxicillin 113 mg PO q12h. Safe dosing for amoxicillin is 80-90 mg/kg/day divided every 12 hours for 5-7 days.**

**Highlight the information that is needed for the nurse to administer the ordered medication safely.**

Amoxicillin is ordered in mg/kg/dose, so it is important to know how much M.S. weighs. Convert pounds to kilograms using the conversion factor: 2.2 pounds = 1 kilogram.

It is important to know how much amoxicillin was ordered for M.S. by the healthcare provider and what the safe dosing range is for M.S. to help determine if the ordered dose is appropriate and safe.

The nurse is preparing to discharge M.S. **Which nursing action is required to ensure the amoxicillin dose is safe for M.S.? (Select all that apply)**

1. Calculate the infant's weight as 2.7 Kg
2. Calculate the infant's weight as 13.2 Kg
3. Calculate the safe dosage range for amoxicillin as: 216-243 mg/day or 108-121.5 mg/dose every 12 hours
4. Calculate the safe dosage range for amoxicillin as 480-540 mg/day or 240-270 mg/dose every 12 hours

1. Calculate M.S.'s weight as 2.7 kg
2. Calculate M.S.'s weight as 13.2 mg
3. Calculate the safe dosage range for amoxicillin as: 216-243 mg/day or 108-121.5 mg/dose every 12 hours
4. Calculate the safe dosage range for amoxicillin as 480-540 mg/day or 240-270 mg/dose every 12 hours

To figure out M.S.'s weight convert pounds to kilograms:

$$\underline{6 \text{ pounds}} \times \underline{1 \text{ kg}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

NGN Unfolding case study: Pediatric calculations and safety

1            2.2 pounds            \*\*the 1 underneath 6 pounds is a place holder and is not significant

The safe dosage range for amoxicillin is 80-90 mg/kg/day:

$$\frac{2.7 \text{ kg}}{1} \times 80 \text{ mg} = 216 \text{ mg}$$

The safe dosage would be between 216-243 mg/day every 12 hours

$$\frac{2.7 \text{ kg} \times 90 \text{ mg}}{1} = 243 \text{ mg}$$

\* divide the safe dosage per day 216-243 by the number of doses being given per day. If a medication is given q12h, then it will be given twice a day [24 hours divided by 12 hours = 2]

$$216 \text{ divided by } 2 = 108$$

A dose between 108-121.5 mg every 12 hours is safe

$$243 \text{ divided by } 2 = 121.5$$

113 mg PO q12h is a safe dose because it is within the safe dosage range of 108-121.5 mg PO q12h. The nurse can comfortably dispense the medication to the parents and show them how much amoxicillin to draw up into the syringe.

-You would get 13.2 kilograms if you mixed up the conversion of pounds to kilograms:

$$\frac{6 \text{ pounds}}{1} \times \frac{2.2 \text{ kg}}{1 \text{ pound}} = 13.2 \text{ kilograms}$$

\*\*this would result in an overdose of any medication that is based on weight. Most medications given to a pediatric patient are based on weight, making it extremely important to understand the appropriate conversion.

The correct conversion is:

$$\frac{6 \text{ pounds}}{1} \times \frac{1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

-480-530 mg/day or 240 -270 mg/dose is incorrect because M.S.'s weight was not converted into kilograms and 6 pounds was used as the weight. This would be equivalent to M.S. weighing a little over 13 pounds. Be careful to make sure the weight used is in kilograms and not pounds. Using the incorrect weight can result in an overdose of medication for pediatric patients.

The safe dosage range for amoxicillin is 80-90 mg/kg/day:

$$\frac{6 \text{ kg}}{1} \times 80 \text{ mg} = 480 \text{ mg}$$

The safe dosage would be between 480-540 mg/day every 12 hours

$$\frac{6 \text{ kg} \times 90 \text{ mg}}{1} = 540 \text{ mg}$$

NGN Unfolding case study: Pediatric calculations and safety

\* divide the safe dosage per day 216-243 by the number of doses being given per day. If a medication is given q12h, then it will be given twice a day [24 hours divided by 12 hours = 2]

480 divided by 2 = 240  
540 divided by 2 = 270

A dose between 240-270 mg is not safe because it is an overdose because an incorrect weight was used

**Choose the *most likely* options for the information missing from the statement below by selecting from the lists of options provided.**

The amoxicillin dose of 113 is \_\_\_\_\_ **1** \_\_\_\_\_ and \_\_\_\_\_ **2** \_\_\_\_\_. The nurse would \_\_\_\_\_ **3** \_\_\_\_\_.

<b>1</b>	<b>2</b>	<b>3</b>
Too high	Safe	Call the physician
Too low	Unsafe	Call the pharmacy
Within range	Lethal	Give the parents the medication

113 mg PO q12h is a safe dose based on M.S.'s weight of 6 pounds, the healthcare provider would not need to be called and the nurse can dispense the medication to the parents after showing them how much amoxicillin to draw up into the syringe.

The correct conversion is

$$\frac{2.7 \text{ kg}}{1 \text{ kg}} \times 80 \text{ mg} = 216 \text{ mg}$$

The safe dosage would be between 216-243 mg/day every 12 hours

$$\frac{2.7 \text{ kg}}{1 \text{ kg}} \times 90 \text{ mg} = 243 \text{ mg}$$

\* divide the safe dosage per day 216-243 by the number of doses being given per day. If a medication is given q12h, then it will be given twice a day [24 hours divided by 12 hours = 2]

216 divided by 2 = 108  
243 divided by 2 = 121.5

A dose between 108-121.5 mg every 12 hours is safe

113 mg PO q12h is a safe dose because it is within the safe dosage range of 108-121.5 mg PO q12h. The nurse can comfortably dispense the medication to the parents and show them how much amoxicillin to draw up into the syringe.

NGN Unfolding case study: Pediatric calculations and safety

1	2	3
Use a 1 mL syringe to draw up 0.13 mL of epinephrine 1 mg/mL solution	Vastus Lateralis (Anterolateral thigh)	264 mL
Use a 3 mL syringe to draw up 0.06 mL of epinephrine 1 mg/mL solution	Gluteus maximus	54 mL
Use a 1 mL syringe to draw up 0.03 mL of epinephrine 1 mg/mL solution	Deltoid	20 mL

-begin by converting M.S.'s weight of 6 pound to kilograms:


$$\frac{6 \text{ pounds} \times 1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

-next figure out the dose of epinephrine to give (1mg/1mL is the standard concentration used for allergic reactions).

$$\frac{0.01 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 0.027 \text{ mL} = 0.03 \text{ (use rounding rules)}$$


\*\* use a 1 mL syringe to draw up the dose because of the small volume, the marking for 0.1 mL and the smaller markings above it indicated 0.01 mL per marking. The smaller markings allow 0.03 mL to be drawn up with accuracy



For an infant, the preferred muscle to use for an intramuscular injection is the vastus lateralis (anterolateral thigh) muscle because it will tolerate the volume the best.



NGN Unfolding case study: Pediatric calculations and safety



M.S.'s weight has already been calculated to be 2.7 kg, calculate the amount of normal saline he would receive as a fluid bolus. The healthcare provider ordered 20 mL/kg of normal saline:

$$\frac{20 \text{ mL} \times 2.7 \text{ kg}}{\text{Kg} \quad 1} = 54 \text{ mL}$$

- give M.S. 54 mL of normal saline over 60 minutes as a fluid bolus once an intravenous line has been started.

Be careful of the conversion from pounds to kilograms. You would get a volume 0.13 mL if you converted pounds to kilograms using the incorrect conversion factor (2.2 kilogram = 1 pound)

$$\frac{6 \text{ pounds}}{1} \times \frac{2.2 \text{ kg}}{1 \text{ pound}} = 13.2 \text{ kilograms}$$

The correct conversion is:

$$\frac{6 \text{ pounds}}{1} \times \frac{1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

This would give you an incorrect dosage of epinephrine:

$$\frac{0.01 \text{ mg}}{1} \times 13.2 \text{ kg} = 0.132 \text{ mL} = 0.13 \text{ (use rounding rules)}$$

NGN Unfolding case study: Pediatric calculations and safety

Kg 1

\*\*this would result in an overdose of epinephrine which can lead to extreme tachycardia and severe hypertension.


The correct dosage is:

$$\frac{0.01 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 0.027 \text{ mL} = 0.03 \text{ (use rounding rules)}$$

A dose of 0.06 mL would be calculated if you did not convert 6 pounds to kilograms. It is important to make sure you have the correct weight in the unit needed. Most medications are dosed in kilograms, so make sure the weight is in kilograms. A kilogram is smaller than a pound, so you can determine if the weight is in the correct units.

$$\frac{0.01 \text{ mg}}{\text{Kg}} \times \frac{6 \text{ kg}}{1} = 0.06 \text{ mL} = \text{(this would result in giving the M.S. twice the prescribed dose)}$$

You would not be able to accurately draw up 0.06 mL in a 3 mL syringe, the markings at 0.1 mL, 0.06 mL would be between the first marking and the end of the syringe, making it almost impossible to accurately draw up the correct dose.



The correct dosage is: 0.03 mL

The gluteus maximus and the deltoid muscle are not developed enough at 6 months of age to be able to absorb an intramuscular injection. The preferred muscle is the vastus lateralis in the anterolateral thigh for children 6 months and under. It is important to make you are giving intramuscular injections into a muscle that is able to handle the volume and be able to appropriately absorb the medication.

-The correct weight and units is important when calculating pediatric dosages.

$$\frac{20 \text{ mL}}{\text{Kg}} \times \frac{13.2 \text{ kg}}{1} = 264 \text{ mL} \quad \text{(you mixed up the conversion and use 2.2 kg = 1 pound)}$$

this would give M.S. significantly more fluid than was ordered and it could lead to pulmonary edema and breathing problems

$$\frac{20 \text{ mL}}{\text{Kg}} \times \frac{6 \text{ kg}}{1} = 120 \text{ mL} \quad \text{(you did not convert 6 pounds to kilograms and used 6 pounds in place of the ordered kilograms). This would give M.S. more fluid than was}$$

NGN Unfolding case study: Pediatric calculations and safety

ordered by the physician. It is important to make sure you have the correct weight in the correct units.

The correct calculation is:

$$\frac{20 \text{ mL} \times 2.7 \text{ kg}}{\text{Kg} \quad 1} = 54 \text{ mL}$$

<b>Actions</b>	<b>Indicated</b>	<b>Contraindicated</b>
Administer Ceftriaxone 300 mg IV		x
Administer Ceftriaxone 135 mg IV	x	
Calculate the safe dosage range for ibuprofen as 13.5-27 mg per dose. The dose of 20 mg PO Q8h PRN is safe and should be administered.	x	
Calculate the safe dosage range for ibuprofen as 66-132 mg per dose. The dose of 20 mg PO q8H PRN is low and the healthcare provider should be notified.		x
Calculate the safe dosage range for acetaminophen as 27-40.5 mg per dose. The dosage of 64 mg PO q4h PRN is too high and the healthcare provider should be notified.	x	
Calculate the safe dosage range for acetaminophen as 60-90 mg per dose. The dosage of 64 mg PO q4h PRN is safe and should be administered.		x

-begin by calculating how many kilograms M.S. weighs:

$$\frac{6 \text{ pounds} \times 1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

The physician ordered Ceftriaxone 50 mg/kg IV daily:

$$\frac{50 \text{ mg} \times 2.7 \text{ kg}}{\text{Kg} \quad 1} = 135 \text{ mg}$$

NGN Unfolding case study: Pediatric calculations and safety

The safe dose for ibuprofen is 5-10 mg/kg per dose PO q6-8h PRN. Calculate the safe dosage range for M.S.

$$\frac{5 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 13.5 \text{ mg}$$

The safe dosage range for ibuprofen would be 13.5-20 mg  
20 mg PO Q8h is within the safe dosage range for M.S. and

$$\frac{10 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 27 \text{ mg}$$

should be administered to help with M.S.'s fever

The safe dose for acetaminophen is 10-15 mg/kg per dose q4-6h PRN. Calculate the safe dosage range for M.S.

$$\frac{10 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 27 \text{ mg}$$

The safe dosage range for acetaminophen would be 27-40.5 mg  
64 mg PO q4h is NOT within the safe dosage range for M.S. and

$$\frac{15 \text{ mg}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 40.5 \text{ mg}$$

should NOT be administered, the s  
healthcare provider should be called to clarify t

the order

Be careful of the conversion from pounds to kilograms. a volume 0.13 mL is calculated when the incorrect conversion factor for pounds to kilograms is used(2.2 kilogram = 1 pound)

$$\frac{6 \text{ pounds}}{1 \text{ pound}} \times \frac{2.2 \text{ kg}}{1} = 13.2 \text{ kilograms}$$

Make sure to convert 6 pounds to kilograms. Pounds and kilograms are not the same unit. The pound is a larger unit than the kilogram. The correct conversion factor is 2.2 pounds = 1 kilogram. If 6 pounds is assumed to be the same a 6 kilograms, the patient is getting at least twice the ordered dose.

The healthcare provider ordered Ceftriaxone 50 mg/kg IV daily:

$$\frac{50 \text{ mg}}{\text{Kg}} \times \frac{6 \text{ kg}}{1} = 300 \text{ mg}$$

(convert 6 pounds to kilograms. Kilograms and pounds are  
not the same unit, be careful to make appropriate  
conversions.)

The correct conversion is:

$$\frac{6 \text{ pounds}}{1} \times \frac{1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

NGN Unfolding case study: Pediatric calculations and safety

The safe dose for ibuprofen is 5-10 mg/kg per dose PO q6-8h PRN. Calculate the safe dosage range for M.S.

$$\frac{6 \text{ pounds}}{1 \text{ pound}} \times 2.2 \text{ kg} = 13.2 \text{ kilograms}$$

(the conversion of pounds to kilogram. 1 kilogram = 2.2 pounds. The kilogram is smaller than the pound)

$$\frac{5 \text{ mg}}{\text{Kg}} \times 13.2 \text{ kg} = 66 \text{ mg}$$

The safe dosage range for ibuprofen is not 66-132 mg  
20 mg PO Q8h is low according to this calculation.

$$\frac{10 \text{ mg}}{\text{Kg}} \times 13.2 \text{ kg} = 132 \text{ mg}$$

This dose would be an overdose

The correct dosage calculation is:

The safe dose for ibuprofen is 5-10 mg/kg per dose PO q6-8h PRN. Calculate the safe dosage range for M.S.

$$\frac{5 \text{ mg}}{\text{Kg}} \times 2.7 \text{ kg} = 13.5 \text{ mg}$$

The safe dosage range for ibuprofen would be 13.5-20 mg  
20 mg PO Q8h is within the safe dosage range for M.S. and

$$\frac{10 \text{ mg}}{\text{Kg}} \times 2.7 \text{ kg} = 27 \text{ mg}$$

should be administered to help with M.S.'s fever

The safe dose for acetaminophen is 10-15 mg/kg per dose q4-6h PRN. Calculate the safe dosage range for M.S.

You did not convert 6 pounds to kilograms. You used 6 pounds as an equivalent to kilograms. The kilogram is a smaller unit than the pound and it is important to make sure correct conversions are completed for pediatric patients.

$$\frac{10 \text{ mg}}{\text{Kg}} \times 6 \text{ kg} = 60 \text{ mg}$$

The safe dosage range for acetaminophen is not 60-90 mg  
64 mg PO q4h is NOT within the safe dosage range for M.S. and

$$\frac{15 \text{ mg}}{\text{Kg}} \times 6 \text{ kg} = 90 \text{ mg}$$

should NOT be administered, the  
healthcare provider should be called to clarify

the order. This would result in an overdose and could lead to liver damage

The safe dose for acetaminophen is 10-15 mg/kg per dose q4-6h PRN. Calculate the safe dosage range for M.S.

$$\frac{10 \text{ mg}}{\text{Kg}} \times 2.7 \text{ kg} = 27 \text{ mg}$$

The safe dosage range for acetaminophen would be 27-40.5 mg  
64 mg PO q4h is NOT within the safe dosage range for M.S. and

$$\frac{15 \text{ mg}}{\text{Kg}} \times 2.7 \text{ kg} = 40.5 \text{ mg}$$

should NOT be administered, the healthcare provider should be  
called to clarify the order

NGN Unfolding case study: Pediatric calculations and safety

Nursing Action	Indicated	Contraindicated	Non-Essential
Give a normal saline fluid bolus of 120 mL		x	
Give a normal saline fluid bolus of 54 mL	x		
Calculate the norepinephrine dose as 1mL/hr	x		
Calculate the norepinephrine dose as 2.3 mL/hr		x	
Give acetaminophen to help with the fever	x		

-begin by calculating how many kilograms M.S. weighs:

$$\frac{6 \text{ pounds}}{2.2 \text{ pounds}} \times \frac{1 \text{ kg}}{2.2 \text{ pounds}} = 2.72 = 2.7 \text{ (using round rules: 4 or less round down and 5 or higher round up)}$$

\*\*the 1 underneath 6 pounds is a place holder and is not significant

The healthcare provider ordered 20 mL/kg of normal saline:

$$\frac{20 \text{ mL}}{\text{Kg}} \times \frac{2.7 \text{ kg}}{1} = 54 \text{ mL}$$

\*you would want to give M.S. 54 mL of normal saline over 60 minutes per the physician order

Norepinephrine is ordered as 0.05 mcg/kg/min. Calculate the dose in mL/hr

$$\frac{2.7 \text{ kg} \times 0.05 \text{ mcg}}{1 \text{ mcg/min}} \times \frac{1 \text{ mg}}{1000 \text{ mcg}} \times \frac{250 \text{ mL}}{2 \text{ mg}} \times \frac{60 \text{ min}}{1 \text{ hour}} = 1.01 \text{ mL/hr} = 1 \text{ mL/hr (rounding rules)}$$

\* start the norepinephrine at 1 mL/hr and monitor M.S. very closely for changes in blood pressure. A patient that requires blood pressure support and vasopressor medications is in an intensive care unit to allow for close monitoring.

Controlling fever is an essential part of managing sepsis, so after blood pressure medication as been started, acetaminophen should be given to help decrease fever.

You need to make sure your patient is breathing, has a heart rate, and an adequate blood pressure before addressing any other issues. The patient needs to be able to sustain life and then you address the non-life threatening issues.

NGN Unfolding case study: Pediatric calculations and safety

$$\frac{20 \text{ mL}}{\text{Kg}} \times \frac{6 \text{ kg}}{1} = 120 \text{ mL}$$

\* give M.S. 120 mL of normal saline over 60 minutes because 6 pounds was not converted to kilograms  
It is very important to make sure correct units are used

Norepinephrine is ordered as 0.05 mcg/kg/min. You need to calculate the dose in mL/hr

$$\frac{6 \text{ kg} \times 0.05 \text{ mcg} \times 1 \text{ mg} \times 250 \text{ mL} \times 60 \text{ min}}{1 \text{ mcg/min} \times 1000 \text{ mcg} \times 2 \text{ mg} \times 1 \text{ hour}} = 2.25 \text{ mL/hr} = 2.3 \text{ mL/hr (rounding rules)}$$

Six pounds was not converted to kilograms, giving the patient too much norepinephrine. Giving too much of the medication could result in a significantly higher blood pressure and a significant increase in heart rate. It is very important to make sure the correct weight and the correct units are used before administering any medication.

The correct calculation is:

$$\frac{2.7 \text{ kg} \times 0.05 \text{ mcg} \times 1 \text{ mg} \times 250 \text{ mL} \times 60 \text{ min}}{1 \text{ mcg/min} \times 1000 \text{ mcg} \times 2 \text{ mg} \times 1 \text{ hour}} = 1.01 \text{ mL/hr} = 1 \text{ mL/hr (rounding rules)}$$

After 5 days in the hospital, M.S. has a temperature of 37 degrees Celsius and is off of the norepinephrine drip. The infant's parents want to take the baby home. The healthcare provider assesses a minor infection remains in the left ear and orders continued antibiotic therapy at home. Orders include azithromycin 25 mg PO daily for 3 days. M.S. currently weighs 5.5 pounds. The safe dose for azithromycin is 10 mg/kg daily for 3 days.

**Highlight the information necessary for the nurse to determine if the azithromycin dose is safe for M.S. before sending him home.**

To determine if a dose is safe for M.S. determine how much M.S. weighs in kilograms.

$$\frac{5.5 \text{ pounds} \times 1 \text{ pound}}{1} = \frac{2.5 \text{ kg}}{2.2 \text{ kg}}$$

-determine if the dose is safe. The safe dose is azithromycin 10 mg/kg daily

$$\frac{2.5 \text{ kg} \times 10 \text{ mg}}{\text{kg}} = 25 \text{ mg}$$

The dose is safe of M.S. and the nurse can teach the parents how to draw up the appropriate amount of azithromycin to give to M.S.

\*\*to determine safe dosages, you need to know what the safe dose is and the patient's weight in the correct units\*\*

**References:**

- Gray Morris, D.C. (2018). *Calculate with confidence* (7<sup>th</sup> ed). St. Louis, MO: Elsevier
- Kee, J., Marshall, S., Forrester, M., & Woods, K. (2021). *Clinical calculations with applications to general and specialty areas* (9<sup>th</sup> ed). St. Louis, MO: Elsevier
- Micromedex (2020)
- Ogden, S.J. & Fluharty, L.K. (2020). *Calculation of drug dosages: a work text* (11<sup>th</sup> ed). St. Louis, MO: Elsevier
- Tritak, A. & Daingerfield, M. (2020). *Brown and Mulholland's drug calculations: Ratio and proportion problems for clinical practice* (11<sup>th</sup> ed). St. Louis, MO: Elsevier
- Turner, S.J. (2019). *Mulholland's the nurse, the math, the meds: Drug calculations using dimensional analysis* (4<sup>th</sup> ed). St. Louis, MO: Elsevier